arXiv:quant-ph/0405136v1 23 May 2004

Coherent manipulation of motional states of trapped ions

Hao-Sheng Zeng[1,2*], Le-Man Kuang[1] and Ke-Lin Gao[2]

[1] *Department of Physics, Hunan Normal University, Hunan 410081, People's Republic of China*

[2] *Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Wuhan Institute of Physics and Mathematics, Chinese Academy of Science, Wuhan 430071, People's Republic of China*

Abstract

The beam splitter and phase shifter, which are the key elements in the experiments of light interference, are realized in the motion of trapped ions. Some applications, such as the creation of quantum motional states and the realization of Mach-Zehnder interferometer, are illustrated. Several detection methods of motional states used in the interferometer are also discussed.

Keywords: Quantum motional states, beam splitter, phase shifter.

Typeset using REVTEX

---

*E-mail adress: hszeng@mail.hunnu.edu.cn

1. Introduction

Beam splitter is a very important element in quantum optical experiments which can realize the splitting and mixing of the input light fields to produce correlated or entangled output states. A number of authors have considered the behavior of the quantum-mechanical beam splitter in the past few years [1-6]. In nearly every interferometric apparatus such as Mach-Zehnder[1, 7-9], Fabry-Perot[8] and Michelson-type[10] interferometers, the beam splitter plays a central role. Both passive and active interferometers [10-12] that have been touted to revolutionize ultrasensitive measurements like gravity-wave detection, sensing of tiny rotation rates, etc. use beam splitters. In addition, phase shifter, which may be realized by changing the length of an optical path or by employing the optical Kerr effect [10, 13], is also an important ingredient in most optical interference experiments.

On the other hand, the developments in the cooling and trapping ions[14] have opened a new research for both atomic physics and quantum optics. When ions trapped in an electromagnetic harmonic potential, the quantized vibratic motions play a role of boson modes. And the interaction between ions and laser fields leads the coupling between the external and internal degrees of freedom of the trapped ions[15-16]. As the coupling between the vibratic modes and the external environment is extremely weak, dissipative effects, which are inevitable from cavity damping in the optical regime, can be significantly suppressed. This unique feature thus can be used to realize some cavity QED experiments without using an optical cavity. Stimulated by these ideas, many schemes have been proposed for generating nonclassical motional states of trapped ions, such as Fock states [16], squeezed state [17-19], even and odd coherent states [20-23] and their various kinds of superpositions[24-29]. These nonclassical motional states may be used to perform quantum computation[30-32], test quantum-mechanical basic principals[32] and study quantum decoherence effects[33]. Up to now, motional Fock states, squeezed states, coherent states [34-35] and Schr$\ddot{o}$dinger cat [36] for the c.m. mode of a single trapped ion have been produced and observed experimentally. The reconstruction of motional states have also been realized in experiments[37].

However, the ability that people coherently manipulate motional states is much lower than that people manipulate light fields. In order to further enrich and develop the content of this portion, we here present a method to implement the beam splitter and phase shifter in the motion of trapped ions, and then use them to coherently produce some typical quantum motional states and to implement the famous Mach-Zehnder interferometer in the vibrational motion of trapped ions. Some of the results or ideas may also be used directly to implement other vibrational interferential experiments, such as the homodyne and Michelson-type interferometer. The construction of this paper is arranged as follows: In section 2, we first describe how to construct the two pivotal elements for implementing a Mach-Zehnder interferometer— the two-mode beam splitter and phase shifter for the motional states of c.m. and breathing modes in two trapped ions, where the later is mainly based on the dynamical evolution of laser-drived trapped ions first introduced by D'Helon and Milburn[38]. In section 3, we then illustrate their applications in the generation of quantum motional states. And in section 4, we further use them to implement a Mach-Zehnder interferometer for the vibrational motion of trapped ions, where three detection ways of motional states are provided. The first two ways are mainly based on the idea of the reconstruction of motional states[34, 39], and the third one based on the direct measurement of the mean value of a motional operator[40]. A conclusion is given in section 5.

2. Beam splitter transformation and phase shift in the motion of trapped ions

Beam splitter and phase shifter are two key elements in quantum optical experiments. Their realizations in the motion of trapped ions will allow us to implement the similar interferential experiments in trapped-ion systems without the use of optical elements. Though some papers have already involved this problem[41-42], the general constructions of beam splitter and phase shifter in trapped-ion systems need further developments. Here we provide a method to realize this two important elements in the motion of two trapped ions.

Following the experimental implementation of reference [43], we consider a two-level system which is formed by two $^9\mathrm{Be}^+$ ions of mass m trapped in a linear trap. The two spectrally resolved ground-state hyperfine levels are defined as the ground and excited states of our problem: $|F=2, m_F=-2\rangle \equiv |g\rangle$ and $|F=1, m_F=-1\rangle \equiv |e\rangle$, which form the so-called internal degrees of freedom. Assume that the two ions are strongly bounded in the y and z directions, but weakly bounded in a harmonic potential in the x direction and oscillate around their equilibrium positions: $x_{10}=-d/2$, $x_{20}=d/2$. We denote by $\widehat{X}=(\widehat{x}_2+\widehat{x}_1)/2$, $\widehat{x}=(\widehat{x}_2-\widehat{x}_1)/2$ the c.m. and breathing mode operators respectively which constitute the studied external vibratic motional degrees of freedom. Now we let only one of the ions, say the first ion, to be illuminated simultaneously by two classical standing wave lasers with the positive frequency parts $E_I^{(+)}=E_{0I}\cos(k_I x+\varphi_I)e^{-i\omega_I t-i\phi_I}$ and $E_{II}^{(+)}=E_{0II}\cos(k_{II}x+\varphi_{II})e^{-i\omega_{II}t-i\phi_{II}}$. Assume that the two lasers have the same amplitudes $E_{0I}=E_{0II}=E_0$ and effective wavevectors $k_I=k_{II}=k$, but different frequencies ω_I and ω_{II}. Experimentally, the two standing wave lasers with needed properties can be implemented by employing two-photon stimulated-Raman transitions[18, 31]. Of course, in order to let a laser illuminate selectively only ion 1 without lighting ion 2, the laser propagation should form an appropriate angle with x axis[44] and the projective wave in the direction of x axis plays the role of coupling the internal and the wanted external degrees of freedom. Assuming that the ion is located at the anti-nodes of both the two standing waves, i.e. $\varphi_I=\varphi_{II}=0$, then the Hamiltonian for this system can be written as,

$$H=H_0+H_{int}, \tag{1}$$

$$H_0=\mu a^+a+\nu b^+b+\omega_0\sigma_{z1}/2, \tag{2}$$

$$H_{int}=\frac{\Omega}{2}\sigma_{+1}[(e^{-i\omega_I t-i\phi_I}+e^{-i\omega_{II}t-i\phi_{II}})\cos(k\widehat{x}_1)+h.c.] \tag{3}$$

where μ (ν) and a (b) are the frequency and annihilation operator of the c.m. mode (breathing mode), ω_0 the energy difference between ground state $|g\rangle$ and excited state $|e\rangle$ of each ion, σ_{z1} and σ_{+1} the usual Pauli operators describing the internal degrees of freedom for ion 1, Ω the Rabi frequency for the ion-laser coupling which is assumed to be equal for the two standing wave lasers.

We select the two standing waves to excite resonantly both the upper and lower sidebands of c.m. and breathing modes, i.e. the frequencies of the two lasers satisfying

$$\omega_I = \omega_0 - (\mu - \nu), \quad \omega_{II} = \omega_0 + (\mu - \nu). \tag{4}$$

Under the condition that both c.m. and breathing modes are cooled to Lamb-Dicke regime, we can expand the Hamiltonian up to the order terms of η^2 or η_r^2 which represent the lowest couplings between internal and external degrees of freedom. Transforming the above Hamiltonian to the interaction picture with respect to H_0, we have

$$H_{int}^1 = \Omega\eta\eta_r(a^+b + ab^+)\sigma_{x1} \tag{5}$$

for one kind of laser-phase selection of $\phi_I = \phi_{II} = 0$ and

$$H_{int}^2 = -i\Omega\eta\eta_r(a^+b - ab^+)\sigma_{x1} \tag{6}$$

for another kind of laser-phase selection of $\phi_I = -\phi_{II} = -\pi/2$, where $\eta = k\sqrt{1/4m\mu}$ and $\eta_r = k\sqrt{1/4m\nu}$ are the Lamb-Dicke parameters corresponding to the c.m. and breathing modes, σ_{x1} the x component of Pauli operator for ion 1. The exact propagators for the above Hamiltonians are

$$U_1 = \exp\{-i\Omega\eta\eta_r t(a^+b + ab^+)\sigma_{x1}\} \tag{7}$$

for the first kind of laser-phase selection and

$$U_2 = \exp\{-\Omega\eta\eta_r t(a^+b - ab^+)\sigma_{x1}\} \tag{8}$$

for the later kind of laser-phase selection. Note that the interactions of U_1 and U_2 occur only between the internal states of ion 1 and motional states of the system. Now we prepare the initial state of the whole system as

$$|\psi\rangle = |\psi\rangle_1 \otimes |\psi\rangle_2 \otimes |\psi\rangle_{c,r} \tag{9}$$

with$|\psi\rangle_1 = (|e\rangle_1 + |g\rangle_1)/\sqrt{2}$ and $|\psi\rangle_2$ (which may be a arbitrary pure state) the internal states of the two ions and $|\psi\rangle_{c,r}$ the initial motional state of c.m. and breathing modes. By performing unitary propagators U_1 or U_2 on this initial state respectively, we find that both the internal states of the two ions remain unchanged but the motional states experiences a beam-splitter transformation:

$$B_1(\theta)\left|\psi\right\rangle_{c,r} = e^{-i\theta J_x}\left|\psi\right\rangle_{c,r}, \tag{10}$$

or

$$B_2(\theta)\left|\psi\right\rangle_{c,r} = e^{-i\theta J_y}\left|\psi\right\rangle_{c,r} \tag{11}$$

respectively, where $B_1(\theta) = \exp(-i\theta J_x)$ and $B_2(\theta) = \exp(-i\theta J_y)$ with $\theta = 2\Omega\eta\eta_r t$ being a real angle. The Schwinger angular-momentum operators [45] are defined as

$$J_x = (a^+b + ab^+)/2,$$

$$J_y = (a^+b - ab^+)/2i,$$

$$J_z = (a^+a - b^+b)/2,$$

which satisfy the usual commutation relations for the Lie algebra of SU(2): $[J_x, J_y] = iJ_z$, $[J_y, J_z] = iJ_x$ and $[J_z, J_x] = iJ_y$. The transformations $B_1(\theta)$ and $B_2(\theta)$ represent rotations by angle θ about $-x$ and $-y$ axes in the angular-momentum space respectively, which would be distinguished as $\pi/2-$ and $\pi-$ coupling beam splitters[1] for c.m. and breathing modes. When $\theta = \pi/2$, they are referred to as 50/50 beam splitters.

The phase shift transformation for motional states of one trapped ion may be realized by changing the trap frequency for a fixed time[46]. Here we construct the phase shifters for c.m. and breathing modes of two trapped ions using the dynamical evolution of laser-drived trapped ions first introduced by D'Helon and Milburn[38]. For convenience in the next sections, we consider an interaction of a classical standing wave with ion 2 in the above two-ion system. Assume that the ion is placed at the node of the standing wave and the detuning $\Delta = \omega_0 - \omega_L$ is set to excite only the c.m. motional degrees of freedom but far away from the corresponding sideband resonance ($|\Delta| \gg \mu$). In Lamb-Dicke limit and for interaction times much greater than the vibrational period of c.m. mode, this interaction produces the following conditional phase shift[38,42]

$$U_3 = \exp[-i\chi t a^+ a(\sigma_{z2} + 1/2)], \tag{12}$$

where $\chi = \eta^2\Omega^2/(2\Delta)$ with η the Lamb-Dicke parameter of c.m. mode and Ω the Rabi frequency for the ion-laser coupling. Pauli operator σ_{z2} describe the internal degrees of freedom for ion 2, and t is the interaction time. The conditional phase shift U_3 occurs only between the internal state of ion 2 and the c.m. motional state which can be used to realize a phase shift transformation on c.m. mode. For example, by preparing the initial state of the involved system as $|g\rangle_2 \otimes |\psi\rangle_c$ with $|g\rangle_2$ the ground state of ion 2 and $|\psi\rangle_c$ the vibrational state of c.m. mode, then the action of U_3 on this state produces a phase shift on c.m. mode:

$$P_c(\varphi)\,|\psi\rangle_c = \exp(i\varphi a^+ a)\,|\psi\rangle_c \tag{13}$$

without changing the internal state of ion 2. The phase-shift angle $\varphi = \chi t/2$ may be adjusted at will by means of altering the interaction time.

The phase shift on breathing mode may also be realized in a similar way. In the above process, if we use the dispersive standing wave to excite non-resonantly the breathing mode instead of c.m. mode, in the similar conditions one has

$$U_4 = \exp[-i\chi_r t b^+ b(\sigma_{z2} + 1/2)], \tag{14}$$

where $\chi_r = \eta_r^2\Omega_r^2/(2\Delta_r)$ with $\eta_r = 3^{-1/4}\eta$ the Lamb-Dicke parameter for breathing mode, Ω_r the Rabi frequency and $\Delta_r = \omega_0 - \omega_L$ the detuning frequency which satisfies $|\Delta_r| \gg \nu$ this time. By preparing the initial state of the involved system as $|g\rangle_2 \otimes |\psi\rangle_r$ with $|\psi\rangle_r$ the vibrational state of breathing mode, then, without changing the internal state of ion 2, the unitary propagator U_4 produces a phase shift for breathing mode:

$$P_r(\varphi_r) = \exp(i\varphi_r b^+ b) \tag{15}$$

with the phase-shift angle $\varphi_r = \chi_r t/2$.

It is worthwhile to point out that the similar two-mode beam splitter transformation of eq.(11) can also be realized in the motion of one ion trapped in a two-dimensional isotropic harmonic potential trap[41]. The merit to use two ions instead of one ion is that it can realizes simultaneously the two kinds of beam-splitter transformations of eqs.(10)-(11) by simply adjusting the initial phases of the applied lasers. Further, combining these two kinds of beam-splitter transformations with the phase shifts given by eqs.(13) and (15), we can construct the most general two-mode beam-splitter transformation[6]. The constructions of the above beam splitter and phase shifter on motional states are simple, which employ at most two standing waves to interact with one ion. Thus, it may be hopeful to demonstrate in current or future experimental conditions.

3. Creation of quantum motional states using beam splitter and phase shifter

As they played roles in the quantum optics, the beam-splitter and phase shifter for motional states constituted in the last section may also possess underlying applications. In this and next sections, we will show this compendiously through concrete examples. In this section, let us first describe the applications in producing quantum motional states. The first kind of important motional states to be produced are the entangled number states. By preparing the initial motional state in eq.(9) as $|\psi\rangle_{c,r} = |1\rangle_c \otimes |0\rangle_r \equiv |1,0\rangle_{c,r}$ with $|1\rangle_c$ and $|0\rangle_r$ the Fock states of c.m. and breathing modes respectively, and performing U_1 or U_2 on it, we obtain the output states for motional states as

$$B_1(\theta)\,|1,0\rangle_{c,r} = \cos(\theta/2)\,|1,0\rangle_{c,r} - i\sin(\theta/2)\,|0,1\rangle_{c,r}\,, \tag{16}$$

$$B_2(\theta)\,|1,0\rangle_{c,r} = \cos(\theta/2)\,|1,0\rangle_{c,r} + \sin(\theta/2)\,|0,1\rangle_{c,r} \tag{17}$$

with $\theta = 2\Omega\eta\eta_r t$ which can be adjusted by altering the interaction time t. When $\theta = \pi/2$, we obtain the maximally entangled single-phonon number states $\left(|1,0\rangle_{c,r} - i\,|0,1\rangle_{c,r}\right)/\sqrt{2}$ and $\left(|1,0\rangle_{c,r} + |0,1\rangle_{c,r}\right)/\sqrt{2}$. Instead, if we prepare the initial motional state as $|\psi\rangle_{c,r} = |1,1\rangle_{c,r}$, then the output states are

$$B_1(\theta)\,|1,1\rangle_{c,r} = \cos\theta\,|1,1\rangle_{c,r} - i\sin\theta\left(|2,0\rangle_{c,r} + |0,2\rangle_{c,r}\right)/\sqrt{2}, \tag{18}$$

$$B_2(\theta)\,|1,1\rangle_{c,r} = \cos\theta\,|1,1\rangle_{c,r} - \sin\theta\left(|2,0\rangle_{c,r} - |0,2\rangle_{c,r}\right)/\sqrt{2}. \tag{19}$$

For $\theta = \pi/2$, the maximally entangled two-phonon number states $\left(|2,0\rangle_{c,r} \pm |0,2\rangle_{c,r}\right)/\sqrt{2}$ have been created. These maximally entangled number states play an important role in quantum measurement theory. Similar motional states have been produced for one ion trapped in a two-dimensional isotropic ion trap[41]. The initial motional number states used above have been already prepared experimentally by simply applying a sequence of Rabi π pulses of laser radiation on the blue sideband, red sideband or carrier [34]. Therefore the method for producing above entangled phonon number states is practical. Once the above beam splitter has been realized in experiment, then the creation of the discussed motional states will become truth.

The second kind of important motional states to be produced are the entangled coherent states. According to the formula for beam splitter transformation

$$B_2(\theta)\,|\alpha\rangle_c\,|\beta\rangle_r = \left|\alpha\cos(\frac{\theta}{2}) - \beta\sin(\frac{\theta}{2})\right\rangle_c \otimes \left|\alpha\sin(\frac{\theta}{2}) + \beta\cos(\frac{\theta}{2})\right\rangle_r \tag{20}$$

with $|\alpha\rangle$ and $|\beta\rangle$ are motional coherent states, if we let $|\psi\rangle_{c,r} = N_\pm[|\alpha\rangle_c \pm |-\alpha\rangle_c] \otimes |0\rangle_r$ with $N_\pm = [2(1 \pm \exp(-2|\alpha|^2)]^{-1/2}$ the normalized constants, then equation (11) leads to a pair of entangled odd and even coherent states:

$$\left|\widetilde{\alpha};\widetilde{\beta}\right\rangle_\pm = N_\pm[|\widetilde{\alpha}\rangle_c\left|\widetilde{\beta}\right\rangle_r \pm |-\widetilde{\alpha}\rangle_c\left|-\widetilde{\beta}\right\rangle_r] \tag{21}$$

with $\widetilde{\alpha} = \alpha\cos(\theta/2)$ and $\widetilde{\beta} = \alpha\sin(\theta/2)$. For a 50/50 beam splitter, i.e. $\theta = \pi/2$, one has $\widetilde{\alpha} = \widetilde{\beta} = \alpha/\sqrt{2}$ and eq.(21) reduces to

$$|\widetilde{\alpha};\widetilde{\alpha}\rangle_\pm = N_\pm[|\widetilde{\alpha}\rangle_c\,|\widetilde{\alpha}\rangle_r \pm |-\widetilde{\alpha}\rangle_c\,|-\widetilde{\alpha}\rangle_r]. \tag{22}$$

The preparing methods for these types of entangled coherent states in the motion of one trapped ion have been provided [22-23, 26]. In the limit of $\widetilde{\alpha} \to \infty$ and $\widetilde{\beta} \to \infty$, the above entangled coherent states become Bell states which can be employed to complete teleportation of continuous variables[47]. In addition, in the initial state of eq.(9), if we let $|\psi\rangle_2 = |g\rangle_2$ and $|\psi\rangle_{c,r} = N_\pm[|\alpha\rangle_c \pm |-\alpha\rangle_c] \otimes |\beta\rangle_r$ (the internal state $|\psi\rangle_1$ of ion 1 may be an arbitrary superposition of $|g\rangle_1$ and $|e\rangle_1$), after performing the transformations of eqs.(8)

and (12) successively, i.e. $U_3(t_3)U_2(t_2)$ with $t_2 = \pi/(4\Omega\eta\eta_r)$ and $t_3 = 2\pi/\chi$, we then obtain another type of entangled coherent states (unnormalized) for motional states

$$\left|\varepsilon_-\right\rangle_c \left|\varepsilon_+\right\rangle_r \pm \left|\varepsilon_+\right\rangle_c \left|\varepsilon_-\right\rangle_r \tag{23}$$

with $\varepsilon_\pm = (\beta \pm \alpha)/\sqrt{2}$ while the internal states of the two ions are immune. The odd two-mode coherent states in eqs. (22) and (23) are maximally entangled states for two-motional modes.

In the generation of the motional entangled coherent states described above, we used the odd and even coherent states as the input states which can be prepared easily by the interaction of a ion with lasers [22-23, 26-27]. The advantages to produce motional states in terms of beam splitter or/and phase shifter are the simplicity and practicability: it can produces very complicated nonclassical motional states through simple networks constituted by them.

4. Mach-Zehnder interferometer for the motional states

The Mach-Zehnder interferometer is a fundamental interferential device which may be applied to the detection of phase sensitivity. In general, the input states of the interferometer are Bose modes which creation and annihilation operators d^+ and d satisfy: $d^+\left|n\right\rangle = \sqrt{n+1}\left|n+1\right\rangle$ and $d\left|n\right\rangle = \sqrt{n}\left|n-1\right\rangle$ with $\left|n\right\rangle$ a number state of the considered Bose mode. Leibfried, et al.[46] have recently simulated a Mach-Zehnder interferential experiment in trapped ion system which used the internal and external degrees of freedom of a trapped ion as the two input Bose modes of the interferometer respectively. The harmonic oscillator mode of the c.m. motion of the ion along one axis in the trap served as one of the Bose modes. In the restricted Hilbert space formed by the two ground-state hyperfine levels $\left|F=2, m_F=-2\right\rangle \equiv \left|1\right\rangle$ and $\left|F=1, m_F=-1\right\rangle \equiv \left|0\right\rangle$ of the ion, the raising and lowering operators σ_+ and σ_- between the levels have the same action as the annihilation and creation operators d and d^+ of a Bose field respectively. Thus the internal spin motion of the trapped ion can simulate another Bose mode in a restricted two-dimensional Hilbert space. Here we implement a Mach-Zehnder interferometer by fully employing two vibrational modes of trapped ions—the c.m. and breathing modes. It need not restrict one of the input states of the interferometer within the two-dimensional number Hilbert space and thus can make more general interferential experiments. The sketch for this interferometer is depicted in Fig.1. It consists of two 50/50 vibrational beam splitters S_1, S_2 and a phase shifter φ on c.m. mode. The output phonon-number distributions for the two modes are detected by D_1 and D_2. Note that the Mach-Zehnder interferometer here works in a temporal rather than a spatial regime. The c.m. and breathing vibrational motions take place in the same spatial region which don't propagate along separate paths in practice, and the introduction of the mirrors in Fig.1 is only for the convenience of illustrating which don't exist actually. In the following, we describe concretely the implement of each element. Now the initial state of the whole system is prepared as

$$|\psi_0\rangle = |\psi_0\rangle_1 \otimes |g\rangle_2 \otimes |in\rangle \tag{24}$$

with $|\psi_0\rangle_1 = (|e\rangle_1 - |g\rangle_1)/\sqrt{2}$ the internal state of ion 1 and $|in\rangle$ the input motional state of c.m. and breathing modes entering the interferometer. At the action of eq.(7) for an interaction time $T = \pi/(4\Omega\eta\eta_r)$, the evolved state is

$$|\psi_1\rangle = |\psi_0\rangle_1 \otimes |g\rangle_2 \otimes e^{i(\pi/2)J_x} |in\rangle \tag{25}$$

In this way, we realize the transformation of motional state for the beam splitter S_1. By performing the unitary transformation of eqs.(12) on this state, we then complete the phase shift φ on c.m. mode and the resulted state becomes

$$|\psi_2\rangle = |\psi_0\rangle_1 \otimes |g\rangle_2 \otimes e^{i\varphi a^+ a} e^{-i(\pi/2)J_x} |in\rangle \tag{26}$$

with $\varphi = \eta^2\Omega^2 t/(4\Delta)$. Performing again the propagator of eq.(7) for the same interaction time T, the final transformation of beam splitter S_2 is completed and the output state becomes as

$$|\psi_3\rangle = |\psi_0\rangle_1 \otimes |g\rangle_2 \otimes |out\rangle\,, \tag{27}$$

with

$$\left|out\right\rangle = e^{i(\pi/2)J_x} e^{i\varphi a^+ a} e^{i(\pi/2)J_x} \left|in\right\rangle \tag{28}$$

the output two-mode motional state. According to the idea of Mach-Zehnder interferometer, we should measure the phonon-number difference in the two modes at the output of the interferometer, or equivalently, the mean value: $\langle J_z \rangle = \langle out | J_z | out \rangle$. Assume that the input motional state is

$$\left|in\right\rangle = \left|0\right\rangle_c \otimes \left|\alpha\right\rangle_r , \tag{29}$$

i.e. the c.m. mode be in a vacuum state $\left|0\right\rangle_c$ and the breathing mode in a coherent state $\left|\alpha\right\rangle_r$(usually with large amplitude $|\alpha|$), then we have

$$\langle J_z \rangle = \frac{1}{2} n \cos\varphi \tag{30}$$

with $n = |\alpha|^2$ the initial phonon number of breathing mode. It is shown obviously that if the mean value of $\langle J_z \rangle$ can be measured by experiment, then the shifted phase φ may be determined. The phase uncertainty for this detection may be estimated. From eqs.(28)-(29), we have

$$\begin{aligned} \left\langle J_z^2 \right\rangle &= \langle out | J_z^2 | out \rangle \\ &= \frac{1}{4} n (1 + n \cos^2\varphi), \end{aligned} \tag{31}$$

and hence

$$(\Delta J_z)^2 = \left\langle J_z^2 \right\rangle - \langle J_z \rangle^2 = \frac{1}{4} n. \tag{32}$$

The phase error is thus given by

$$\Delta\varphi = \frac{\Delta J_z}{\left|\partial \langle J_z \rangle / \partial \varphi\right|} = \frac{1}{\sqrt{n} \sin\varphi}. \tag{33}$$

When $\varphi = \pi/2$, the phase error is limited to the minimum $1/\sqrt{n}$ which is reverse to the square root of the input phonon number. Of course, the sensitivity of an interferometer can be improved by designing the input states appropriately [1].

The function of the detectors D_1 and D_2 is to measure the phonon number difference of the output two motional modes, i.e. $\langle J_z \rangle$. In trapped ion system, the measurement of motional states is usually be realized by first mapping these external degrees of freedom onto the internals and then making a measurement on the internal states. The measurement of internal states may be finished by the method of electron shelving which has the detection efficiency nearly close to one[48]. Below, we present three methods to measure $\langle J_z \rangle$. The first way is to measure the phonon number distributions of the two output modes independently, which may be performed in principle by use of the dynamics of Jaynes-Cummings model [34, 39]. A more general measurements of motional states including the determination of non-diagonal elements of a density matrix may be realized by the method of quantum-state reconstruction [37]. In the above two-ion system, if we let ion 2 interact with a standing wave with the ion located in the node of the standing wave, we can then get the Hamiltonian as

$$H = \omega_0 \sigma_{z2}/2 + \mu a^+ a + \nu b^+ b + H_{int}, \tag{34}$$

where $H_{int} = (\Omega/2)\sigma_{+2} \exp(-i\omega_L t) \sin(k\widehat{x}_2) + h.c.$ with k (ω_L) the effective wavevector (frequency) of the standing wave and $\widehat{x}_2$ (σ_{+2}) the position operator (internal raising operator) of ion 2. The meanings of the other symbols are defined as before. By adjusting the laser frequency to be resonant with the first red sideband of the c.m. mode, i.e. $\omega_L = \omega_0 - \mu$, in Lamb-Dicke limit and under rotating wave approximation, the interaction Hamiltonian in the interaction picture can be reduced as the standard JCM

$$H_{int} = \lambda(\sigma_{+2} a + \sigma_{-2} a^+). \tag{35}$$

with $\lambda = \eta\Omega/2$ the coupling strength. For convenience of narration, we expand the output motional state of eq.(28) as

$$|out\rangle = \sum_{m,n} c_{mn} \left|m\right\rangle_c \left|n\right\rangle_r \tag{36}$$

with $\left|m\right\rangle_c$, $\left|n\right\rangle_r$ the motional Fock states of c.m. and breathing modes respectively and c_{mn} the complex numbers. In general, $|out\rangle$ is a two-mode entangled state. Performing the JCM interaction of eq.(35) on the interferometer output state of eq.(27) for a interaction time τ, followed by a measurement of the internal state of ion 2, then the probability that the ion 2 in the ground state can be expressed as

$$P_g(\tau) = \frac{1}{2}[1 + \sum_m p_m \cos(2\lambda\tau\sqrt{m})] \tag{37}$$

with $p_m = \sum_n |c_{mn}|^2$ the phonon number distribution of c.m. mode. A Fourier transformation of the measured signal $P_g(\tau)$ allows one to determine the distribution probability p_m of the output c.m. mode[34, 39]. The phonon-number distribution probability $p_n = \sum_m |c_{mn}|^2$ for breathing mode can be obtained in a similar way. After obtained the phonon-number distribution probabilities p_m and p_n of the two output modes, we can then easily determine the mean value of J_z as

$$\langle J_z \rangle = \frac{1}{2}(\sum_m m p_m - \sum_n n p_n). \tag{38}$$

The second way to determine $\langle J_z \rangle$ is to measure the expansion coefficient c_{mn}, or equivalently the joint distribution probability $p_{mn} = |c_{mn}|^2$. This can be accomplished in terms of the dynamics of two-modes JCM. In the above laser-ion interaction, if we let the ion 2 locate at the anti-node instead of node of the standing wave, and adjust the frequency of the standing wave to excite simultaneously the both lower sidebands of the two motional modes, i.e. $\omega_L = \omega_0 - (\mu + \nu)$, in Lamb-Dicke limit and under rotating wave approximation, the interaction Hamiltonian in interaction picture is

$$H'_{int} = g(ab\sigma_{+2} + a^+ b^+ \sigma_{-2}) \tag{39}$$

with $g = \Omega\eta\eta_r$ the coupling strength. This kind of two-mode JCM interaction has been studied extensively in quantum optics [49]. Now if we let eq.(27) as the initial state of this JCM, then the dynamical evolution, after inserting eq.(36), leads to

$$|\psi(t)\rangle = \sum_{m,n} c_{mn}[\cos(gt\sqrt{mn})\,|m, n, g\rangle - i\sin(gt\sqrt{mn})\,|m-1, n-1, e\rangle],$$

where the first, second and third marks in Dirac symbols denote the states of c.m. mode, breathing mode and the internal state of ion 2 respectively. The unchanged internal state of ion 1 is eliminated. Now, if we again measure the internal state of ion 2, then the measured probability in ground state is

$$P_g(t) = \frac{1}{2}[1 + \sum_{m,n} p_{mn} \cos(2gt\sqrt{mn})]. \tag{40}$$

In a similar way, we can determine the joint probability distribution p_{mn} of the motional state $|m,n\rangle_{c,r}$ by Fourier transforming the measured signal $P_g(t)$ and then obtain the mean value of J_z in terms of eq.(38) as well as $p_m = \sum_n p_{mn}$ and $p_n = \sum_m p_{mn}$.

The third way, which is also the simplest way, to determine $\langle J_z \rangle$ is the direct measurement of the mean value of a motional operator[40]. In order to do this, we first perform a $\pi/2$-pulse carrier transition[48] on ion 2 to reset the output state of the interferometer of eq.(27) as

$$|\psi_4\rangle = |\psi_0\rangle_1 \otimes |\psi_0\rangle_2 \otimes |out\rangle \tag{41}$$

with $|\psi_0\rangle_2 = (|g\rangle_2 + i\,|e\rangle_2)/\sqrt{2}$ the internal state of ion 2. Then we execute the unitary transformation U_3 of eq.(12) on this state. At this moment, the mean value of $\langle \sigma_{x2} \rangle$ in the final state is

$$\begin{aligned} \langle \sigma_{x2} \rangle &= \langle \psi_4 | \, U_3^{+} \sigma_{x2} U_3 \, | \psi_4 \rangle \\ &= - \langle out | \sin(2\chi t a^{+} a) \, |out\rangle \, . \end{aligned} \tag{42}$$

For small interaction time and not very large number of phonons, we can linearize the operator as[40] $\sin(2\chi t a^{+}a) \simeq 2\chi t a^{+} a$, then the above equation reduces to

$$\langle \sigma_{x2} \rangle = -2\chi t \, \langle out | \, a^{+} a \, |out\rangle \, . \tag{43}$$

In trapped ion system, the measurement of $\langle \sigma_{x2} \rangle$ is very simple which may be completed by applying a single qubit rotation to transform the internal state of ion 2 from σ_z-picture to σ_x-picture, followed by a detection of the ion's fluorescence[48]. Therefore the mean number of phonons of the c.m. mode in the output state of the interferometer $\langle a^{+} a \rangle$ can be evaluated easily from eq.(43). The mean number of phonons of breathing mode $\langle b^{+} b \rangle$ can also be evaluated in a similar way. Thus the mean value of $\langle J_z \rangle = [\ \langle a^+ a \rangle - \langle b^+ b \rangle]/2$ could be obtained.

5. Conclusion

In conclusion, we have proposed a method to realize the transformations of beam splitter and phase shifter in the motion of trapped ions and discussed their applications in the generation of quantum motional states and in the Mach-Zehnder interferometer. Several detection methods for the output motional state of the interferometer have also been discussed. Further, our research works may be directly applied to implement some other interferential experiments of motional states, such as the Homodyne detection and Michelson-type interferometer [10]. Though the implementation process involves some complicated ion-laser interactions, some pivotal techniques such as the basic manipulations of ion-laser interaction [34-36,43,48] and the reconstruction of motional states[37] have been already realized experimentally. We expect that, with the increasing of the experimental precision, our proposals will become truth and produce an important effects in the future coherent control of quantum motional states of trapped ions.

Acknowledgments

This work was supported by the National Fundamental Research Program Grant No. 2001CB309310, the National Natural Science Foundation of China under Grant Nos.10347128, 10325523, 90203018 and 19734006, the EYTF of the Education Ministry of China, STF of Hunan Province, and the Scientific Research Fund of Hunan Provincial Education Department.

REFERENCES

[1] Yurke B, McCall S L and Klauder J R 1986 *Phys. Rev.* A **33** 4033

[2] Prasad S, Scully M O and Martienssen W 1987 *Opt. Commun.* **62** 139

[3] Ou Z Y, Hong C K and Mandel L 1987 *Opt. Commun.* **63** 118

[4] Fearn H and Loudon R 1987 *Opt. Commun.* **64** 485

Fearn H and Loudon R 1989 *J. Opt. Soc. Am.* B **6** 917

[5] Huttner B and Ben-Aryeh Y 1988 *Phys. Rev.* A **38** 204

[6] Campos R A, Saleh B E A and Teich M C 1989 *Phys. Rev.* A **40** 1371

[7] Born M and Wolf E 1975 *Principles of Optics* (Oxford: Pergamon)

[8] Steel W H 1983 *Interferometry*, 2nd ed. (Cambridge: Cambridge University)

[9] Caves C M 1981 *Phys. Rev.* D **23** 1693

Bondurant R S and Shapiro J H 1984 *Phys. Rev.* D **30** 2548

[10] Scully M O and Zubairy M S 1997 *Quantum Optics* (London: Cambridge Press)

[11] Billing H et al. 1979 *J. Phys.* E **12** 1043

Scully M O 1985 *Phys. Rev. Lett.* **55** 2802

[12] Weiss R *et al* 1979 *Sources of gravitational radiation* (Cambridge: Cambridge University Press)

[13] Imoto N, Haus H A and Yamamoto Y 1985 *Phys. Rev.* A **32** 2287

[14] Itano W M and Wineland D J 1982 *Phys. Rev.* A **25** 35

Wineland D J *et al* 1987 *Phys. Rev.* A **36** 2220

Diedrich F *et al* 1989 *Phys. Rev. Lett.* **62** 403

Monroe C *et al* 1995 *Phys. Rev. Lett.* **75** 4011

[15] Cirac J I *et al* 1994 *Phys. Rev.* A **49** 1202

D'Helon C and Milburn G J 1995 *Phys. Rev.* A **52** 4755

Wu Y and Yang X X 1997 *Phys. Rev. Lett.* **78** 3086

[16] Cirac J I *et al* 1993 *Phys. Rev. Lett.* **70** 762

Solano E *et al* 2000 *Phys. Rev.* A **62** 021401

[17] Cirac J I *et al* 1993 *Phys. Rev. Lett.* **70** 556

[18] Heinzen D J and Wineland D J 1990 *Phys. Rev.* A **42** 2977

[19] Zeng H S, Kuang L M and Gao K L 2002 *Phys. Lett.* A **300** 427

[20] de Matos Filho R L and Vogel W 1996 *Phys. Rev. Lett.***76** 608

[21] Poyatos J F *et al* 1996 *Phys. Rev.* A **54** 1532

[22] Gerry C C 1997 *Phys. Rev.* A **55** 2478

[23] Zheng S B 1998 *Phys. Rev.* A **58** 761

[24] Gardiner S A, *et al* 1997 *Phys. Rev.* A **55** 1683

Zheng S B 1998 *Phys. Lett.* A **248** 25

[25] Zheng S B *et al* 2000 *Phys. Rev.* A **62** 033807

Zheng S B and Guo G C 1998 *Eur. Phys. J.* D **1** 107

[26] Solano E *et al* 2001 *Phys. Rev. Lett.* **87** 060402

Solano E *et al* quant-ph/0202072; quant-ph/0101056.

[27] Moya-Cessa H et al. 1999 *Phys. Rev.* A **59** 2920

Zeng H S *et al* 2001 *Commun. Theor. Phys.* **36** 335

[28] Gou S C, Steinbach J and Knight P L 1997 *Phys. Rev.* A **55** 3719

Gou S C, Steinbach J and Knight P L 1996 *Phys. Rev.* A **54** 1014

Gerry C C, Gou S C and Steinbach J 1997 *Phys. Rev.* A **55** 630

[29] Zou X B, Pahlke K and Mathis W 2002 quant-ph/0204070

Zou X B, Pahlke K and Mathis W 2001 quant-ph/0108117

[30] Gulde S *et al* 2003 *Nature* **421** 48

[31] Monroe C *et al* 1995 *Phys. Rev. Lett.* **75** 4714

[32] Munro W J, Milburn G J and Sanders B C 2000 *Phys. Rev.* A **62** 052108

[33] Myatt C J *et al* 2000 *Nature* **403** 269

Turchette Q A *et al* 2000 *Phys. Rev.* A. **62** 053807

[34] Meekhof D A *et al* 1996 *Phys. Rev. Lett.* **76** 1796

[35] Ben-Kish A *et al* 2002 *Phys. Rev. Lett.* **89** 267901

[36] Monroe C *et al* 1996 *Science* **272** 1131

[37] Leibfried D *et al* 1996 *Phys. Rev. Lett.* **77** 4281

[38] D'Helon C and Milburn G J 1996 *Phys. Rev.* A **54** 5141

[39] Leibfried D *et al* 2003 *Rev. Mod. Phys.* **75** 281

[40] Militello B, Messina A and Napoli A 2002 quant-ph/0203111; quant-ph/0205153

[41] Gou S C and Knight P L 1996 *Phys. Rev.* A **54** 1682

[42] Schneider S, James D F V and Milburn G J 2000 *J. Mod. Opt.* **47** 499

Schneider S, James D F V and Milburn G J 1998 quant-ph/9808012

[43] Sackett C A *et al* 2000 *Nature* **404** 256

[44] Cirac J I and Zoller P 1996 *Phys. Rev. Lett.* **74** 4091

[45] Schwinger J 1965 in *Quantum Theory of Angular Momentum*, edited by Beidenharn L

and van Dam H (New York: Academic)

[46] Leibfried D *et al* 2002 *Phys. Rev. Lett.* **89** 247901

[47] Ralph T C *et al* 2003 quant-ph/0306004

van Enk S J and Hirota O 2001 *Phys. Rev.* A **64** 022313

[48] Kiellinski D *et al* 2001 *Science* **291** 1013

Rowe M A *et al* 2001 *Nature* **409** 791

[49] Abdel-Hafez A M 1992 *Phys. Rev.* A **45** 6610

Gou S C 1989 *Phys. Rev.* A **40** 5116

Parkins A S 1990 *Phys. Rev.* A **42** 6873

Joshi A and Puri R R 1990 *Phys. Rev.* A **42** 4336

Fig.1. Principle of the Mach-Zehnder interferometer. The c.m. and breathing modes a and b of the two trapped ions are mixed on the first beam splitter S_1. After a phase shift φ on the c.m. mode, they converge at the second beam splitter S_2. Finally the two output motional states are measured by detectors D_1 and D_2.



D1
aout
S2
bout
D2
b'
ain
S1
a'
φ
bin